

A Tutorial in Connectome Analysis (II): Topological and Spatial Features of Brain Networks

Dr Marcus Kaiser

School of Computing
Science /

Institute of Neuroscience
Newcastle University
United Kingdom

WCU Dept of Brain & Cognitive
Sciences

Seoul National University

South Korea


<http://bcs.snu.ac.kr/>

<http://www.biological-networks.org>


Outline

- Spatial neural networks
- Connection length distributions
- Component placement optimization
- Role of long-distance connections
- Deficits due to changes in long-distance connectivity
(Alzheimer's disease and IQ)

Macaque (rhesus monkey) cortical network




C. elegans neural network







Global level (277 neurons)







Local level (131 neurons)




Most projections connect adjacent neurons (Gamma function for connection length distribution)




Also true for human structural and functional connectivity



Why is there an exponential tail?



X: number of steps until another neuron
is in the range of the axonal growth cone
p: probability that unit space contains a neuron
 $q = 1 - p$




$$P(X = n) = p * q^{n-1}$$

-> exponential distribution


Reducing neural wiring costs

- Minimizing total wire length reduces metabolic costs for connection establishment and signal propagation
- *Component Placement Optimization, CPO*
Every alternative arrangement of network nodes will lead to a higher total wiring length




- Not the case! Reductions by 30-50% possible


More long-distance projections than expected




When long-distance fibers are missing (minimal):



original



minimal




Long-distance connections are short-cuts that help to reduce the characteristic path length (or average shortest path, ASP):
Less long-distance connections -> longer path lengths

Low path lengths = few intermediate processing steps are beneficial


- **Synchrony of near and distant regions**




- **Reduced transmission delays**



- **Less (cross-modal) interference**




- **Higher reliability of transmission**



Linking structure and function

Small-world properties




EEG
synchronization
Network
(functional connectivity)

in Alzheimer's patients
and control group

Alzheimer: Path length vs. task performance

Mini Mental State Examination
(attention, memory, language)




Diamonds:
Alzheimer patients

Empty squares:
Control


Cognition: Path length vs. IQ

Resting state fMRI in 19 subjects (functional connectivity based on coherence)



Correlations between local path length λ (shortest path length from one node to any other node) and IQ (Intelligence Quotient) for medial prefrontal cortex, bilateral inferior parietal cortex and precuneus / posterior cingulate regions ($p < 0.05$, corrected for age).

Summary



4. Spatial properties:

- Preference for connections to neighbours
- Fast processing due to long-distance connections

5. Linking structure and function:

- Alzheimer's disease: path lengths
- IQ: path lengths